


Vitamin A and E circulate in different cattle plasma fractions*L. Hymøller and S.K. Jensen**Aarhus University, Animal Science, Blichersallé 20, 8830 Tjele, Denmark; lone.hymoller@agrsci.dk*

Analysis of fat soluble vitamins in plasma fractions of protein, chylomicrons, and lipoproteins can give information about their plasma transport and physiological function. Vitamin A (A) and vitamin E (E) are known to circulate in different plasma fractions; hence they were used to verify a simple plasma fractionation method for cattle plasma. Blood was collected in Na-EDTA coated tubes from 3 Holstein bull calves, which for 3 weeks prior to sampling were fed 3 kg/d of concentrate. The concentrate contained 3 µg/kg A and 120 mg/kg E. Plasma was isolated by centrifugation at 1,500×g for 10 min. Chylomicrons were floated by ultracentrifugation of 6 ml plasma at 100,000×g for 1 h in a Beckman-Coulter 70.1 fixed angle rotor. Fractions of lipoproteins (LPF) and protein were separated by adding 1.5 ml OptiPrep™ (Axis-Shield, Norway), layering Hepes buffered saline 0.85%, pH 7.4 on top, and applying ultracentrifugation at 300,000×g for 18 h. Fractions were harvested top-down with Pasteur pipette and analysed for A and E by HPLC. Average amounts of A and E found in each plasma fraction, in % of the total amount of A and E in all fractions, were: Protein: A (54%) and E (7%). Chylomicrons: A (13%) and E (15%). LPF1 (VLDL): A (1%) and E (2%). LPF2 (LDL/HDL): A (5%) and E (62%). LPF3 (HDL): A (27%) and E (15%). A is transported in chylomicrons from its site of uptake in the intestines but circulates in plasma bound to specific binding proteins, which are responsible for its biological action in the body. Likewise E is transported from the intestines to the liver in chylomicrons. Binding proteins specific to E is known to exist, however E is found in the LDL/HDL fraction of plasma. This indicates that cattle plasma was separated into relevant plasma fractions using the presented ultracentrifugation method. In conclusion, by fractionating cattle plasma by ultracentrifugation it was shown that A mainly circulated in heavier plasma fractions i.e. protein and HDL, whereas most E was found in fractions representing LDL/HDL.

In different worlds: transport of dietary vitamin D₂ and vitamin D₃ in cattle plasma fractions*L. Hymøller and S.K. Jensen**Aarhus University, Animal Science, Blichersallé 20, 8800 Tjele, Denmark; lone.hymoller@agrsci.dk*

It is often stated that vitamin D₂ (D₂) and D₃ (D₃) have similar physiological effects but studies in cattle showed, that D₂ is less efficient than D₃ at securing vitamin D status, measured as liver derived 25-(OH)-D₂ (25OHD₂) and 25-(OH)-D₃ (25OHD₃). The aim was to investigate if binding to different plasma fractions could explain the physiological inefficiency of D₂ compared to D₃ in cattle. 3 Holstein bull calves were fed 75 µg/d D₃ and had ad lib access to hay, naturally containing D₂. Blood was collected weekly for 11 weeks and plasma isolated by centrifugation. Plasma was fractionated by ultracentrifugation and fractions of protein, chylomicron, and lipoprotein (LPF) analysed for content of 25OHD₂, 25OHD₃, D₂, and D₃. Contents of 25OHD₂ vs 25OHD₃ and of D₂ vs D₃ in each fraction were compared as average of all samples by Student's t-test. Average percentages of each metabolite in each fraction were: Protein: 25OHD₂ (8%) vs 25OHD₃ (75%; P<0.001) and D₂ (10%) vs D₃ (16%; P+0.03). Chylomicron: 25OHD₂ (39%) vs 25OHD₃ (17%; P<0.01) and D₂ (0%) vs D₃ (6%; P<0.01). LPF1 (VLDL): 25OHD₂ (0%) vs 25OHD₃ (0%) and D₂ (52%) vs D₃ (33%; P+0.07). LPF2 (LDL/HDL): 25OHD₂ (9%) vs 25OHD₃ (2%; P+0.07) and D₂ (4%) vs D₃ (31%; P<0.01). LPF3 (HDL): 25OHD₂ (34%) vs 25OHD₃ (6%; P<0.001) and D₂ (0%) vs D₃ (15%; P<0.01). D₂ and D₃ were only transported evenly distributed in protein, whereas the % of D₂ and D₃ found in other fractions differed. If this means that D₂ is transported in more volatile plasma fractions than D₃, then D₂ could be more prone to degradation and excretion, contributing to its lower physiological efficiency. 25OHD₃ was mainly transported in protein whereas 25OHD₂ was transported in chylomicron and LDL/HDL. Associating with designated binding proteins is vital to the function of 25OHD₃. Hence transportation in other plasma fractions than protein could explain a compromised physiological function of 25OHD₂ compared to 25OHD₃. In conclusion, D₂ and 25OHD₂ are transported in different plasma fractions than D₃ and 25OHD₃, respectively, which could offer an explanation to a lower physiological efficiency of D₂ than D₃ in cattle.

Book of Abstracts of the 65th Annual Meeting of the European Federation of Animal Science



**Book of abstracts No. 20 (2014)
Copenhagen, Denmark
25 - 29 August 2014**

Book of Abstracts of the 65th Annual Meeting of the European Federation of Animal Science

Copenhagen, Denmark, 25-29 August, 2014



EAAP Scientific Committee:

H. Simianer
G. Van Duinkerken
H. Spoolder
M. Vestergaard
A. Bernues
M. Klopčič
L. Bodin
C. Lauridsen
N. Miraglia
G. Pollott

Proceedings publication and Abstract Submission System (OASES) by



**Wageningen Academic
Publishers**

EAN: 9789086862481
e-EAN: 9789086867998
ISBN: 978-90-8686-248-1
e-ISBN: 978-90-8686-799-8
DOI: 10.3920/978-90-8686-799-8

ISSN 1382-6077

First published, 2014

© Wageningen Academic Publishers
The Netherlands, 2014

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned. Nothing from this publication may be translated, reproduced, stored in a computerised system or published in any form or in any manner, including electronic, mechanical, reprographic or photographic, without prior written permission from the publisher:
Wageningen Academic Publishers
P.O. Box 220
6700 AE Wageningen
The Netherlands
www.WageningenAcademic.com
copyright@WageningenAcademic.com

The individual contributions in this publication and any liabilities arising from them remain the responsibility of the authors.

The designations employed and the presentation of material in this publication do not imply the expression of any opinion whatsoever on the part of the European Federation of Animal Science concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

The publisher is not responsible for possible damages, which could be a result of content derived from this publication.